

Current Developments in our Understanding of Auditory Neuropathy ANSD

[Auditory Neuropathy Spectrum Disorder]

Jon K. Shallop, Ph.D.
Professor Emeritus
Mayo Clinic Rochester, MN

Institut Raymond-Dewar Montreal, Canada June 25, 2009

BACKGROUND INFORMATION

- **Starr et al. (1991) identified auditory neuropathy (AN) as a timing disorder**
- **Zeng et al. (1999) demonstrated temporal processing problems in adults with auditory neuropathy.**

<http://www.com.uci.edu/hesp/home.html>

NeuroReport 10, 3429-3435 (1999)

Temporal and speech processing deficits in auditory neuropathy

Fan-Gang Zeng,^{CA} Sandy Oba,¹ Smita Garde,² Yvonne Sininger¹ and Arnold Starr³

Department of Hearing and Speech Sciences and Center for Comparative Neuroscience, 0100 Lefrak Hall, University of Maryland, College Park, MD 20742; ¹House Ear Institute, Los Angeles, CA 90057; ²University of Southern California, Los Angeles, CA 90089; ³Department of Neurology, University of California, Irvine, CA 92717, USA


AUDITORY neuropathy affects the normal synchronous activity in the auditory nerve, without affecting the amplification function in the inner ear. Patients with auditory neuropathy often complain that they can hear sounds, but cannot understand speech. Here we report psychophysical tests indicating that these patients' poor speech recognition is due to a severe impairment in their temporal processing abilities. We also simulate this temporal processing impairment in normally hearing listeners and produce similar speech recognition deficits. This study demonstrates the importance of neural synchrony for auditory perceptions including speech recognition in humans. The results should contribute to better diagnosis and treatment of auditory neuropathy. *NeuroReport* 10:3429-3435 © 1999 Lippincott Williams & Wilkins.

Key words: Acoustic simulation; Auditory neuropathy; Hearing disorder; Human; Neural synchronization; Speech perception; Temporal processing

^{CA}Corresponding Author

The characteristics of Hearing Loss in ANSD

Zeng et al. *NeuroReport* 10, 3429-3435 (1999)



BACKGROUND INFORMATION

- [Lesinski-Schiedat A.](#) [Frohne C.](#) [Hemmaouil I.](#) [Battmer RD.](#) [Lenarz T.](#)
Funktionelle Taubheit bei peri-synaptischer Audiopathie - Isolierte Storungen der inneren Haarzellen? Laryngo- rhino-otologie. 80(10):601-4, 2001
- Subjective deafness in case of **peri-synaptic audiopathy**. Isolated defects of the inner haircells?
- "Hypoxia, carboplatin, ototoxicity and metabolic disorders are possible etiologies for damage to the inner hair cells or synapsis."

5

BACKGROUND INFORMATION

- [Rodriguez-Ballesteros M.](#) [del Castillo FJ.](#) [Martin Y.](#) [Moreno-Pelayo MA.](#) [Morera C.](#) [Prieto F.](#) [Marco J.](#) [Morant A.](#) [Gallo-Teran J.](#) [Morales-Angulo C.](#) [Navas C.](#) [Trinidad G.](#) [Tapia MC.](#) [Moreno F.](#) [del Castillo I.](#)
- Unidad de Genetica Molecular, Hospital Ramon y Cajal, Madrid, Spain.
- Auditory neuropathy in patients carrying mutations in the **otoferlin gene** (OTOF).
- ***Human Mutation***. 22(6):451-6, 2003 Dec.

Varga R, Kelley PM, Keats BJ, et al. Non-syndromic Recessive Auditory Neuropathy is the Result of Mutations in the **Otoferlin (OTOF) Gene. *J Med Genet* 40:45-50, 2003.**

BACKGROUND INFORMATION

- Berlin et al. (2001), and Hood (2002) proposed auditory dys-synchrony as a more descriptive term. Typical findings include:
 - abnormal evoked potentials
 - otoacoustic emissions (usually present)
 - absent middle ear reflexes
 - lack of benefit from amplification

7

Prevalence

There are no known data on how many children with ANSD have been implanted but if we assume that there are approximately **140,000** cochlear implants worldwide at the present time (Wilson and Dorman, 2007) and that **40%** of them are children, then there are at least **56,000** children worldwide with cochlear implants.

8

Prevalence

A conservative estimate of **10%** for ANSD (Uus and Bamford, 2005) can be argued which suggests that more than **5,000** children with cochlear implants worldwide are likely to have ANSD as a component of their hearing loss etiology.

9

Table 1: Prevalence of ANSD in “at-risk” infant populations

Study	Population	# Subjects	No. of ANSD	subjects % of Total
Stein et al. (1996)	Special Care Nursery	100	4	4.00
Psarommatis et al. (1997)	Intensive Care Unit	102	2	1.96
Rance et al. (1999)	“at-risk” infants	5199	12	0.23

Rance 2005 AN/AD and it's Perceptual Consequences

10

Table 2: Prevalence of ANSD in children with permanent hearing loss.

Study	Population	No. of Cases Permanent Hg. Loss	No. of AN/AD Cases	% of Total
Kraus et al. (1984)	Hg. impaired children	48	7	14.58
Park & Lee (1998)	Hg. impaired children	139	7	5.04
Vohr et al. (1998)	Universal screening	111	2	1.80
Rance et al. (1999)	“at-risk” infants	109	12	11.01
Berlin et al.	Hg. impaired	1000	87	8.70

Rance 2005 AN/AD and it's Perceptual Consequences

The characteristics and co-morbidities of ANSD

Children with ANSD can present with some unique characteristics that may initially be misunderstood by parents teachers and clinicians.

This also implies that some children presently in classrooms may in fact have this disorder.



The characteristics and co-morbidities of ANSD

In general, various physical, sensory or cognitive issues can be observed in addition to hearing loss and all of these will be educationally significant.

As these “issues” arise, there is an increase in the probability of auditory neuropathy [ANSD] as an additional problem.

13



The characteristics and co-morbidities of ANSD

- Developmental delays
- Learning disabilities, ADD, ADHD
- Autism spectrum disorders
- Emotional or behavioral problems
- Uncorrected visual problems, blindness

14



The characteristics and co-morbidities of ANSD

- **Other health impairment**
- **Cerebral palsy, motor disorders**
- **Apraxia**
- **Inner ear malformations, Atretic or absent auditory nerve**
- **Seizures**
- **Syndromes**
- **Other**

15

Lessons Learned:


The characteristics and co-morbidities of ANSD

- **Hearing is usually abnormal in these children**
- **However they may exhibit good sound "detection"**
- **And very poor speech recognition in most cases**
- **But some will benefit from hearing aids**
- **.... And why ? Sound demonstration**

16

Acoustic Simulations of ANSD

Notice the acoustic 'smearing'




Simulation : from “profound” to “original”
speech segments progressively overlap
resulting in a “smearing” effect.

Fan-Gang Zeng et al. (1999)

17

Gap Detection

Speech is a series of
sequential events with
“gaps” between words



18

Gap Detection

However, what if the words overlap ?



mamaison est votre maison

19

Gap Detection

What is gap detection and why is it important for speech recognition ?



20

Gap Detection


What is gap detection and why is it important for speech recognition ?

Speech is segmented



21


Gap Detection is abnormal for persons with ANSD



Zeng et al 1998

22

Gap Detection is abnormal for persons with ANSD



Zeng et al 1998

23

Lessons Learned

Speech recognition is very dependant on the ability to make fine timing discriminations in the speech signal which has small temporal gaps and transitions

24

Gap Detection

Some examples of the importance of gap detection

“cue” vs “coo” /ku/ vs. /kiu/



25

Gap Detection

Some examples of the importance of gap detection

“stop” vs “shop”



26


Case Study #1

Profound Hearing Loss in the Right ear ?

- * Passed OAE screening in both ears
- * Absent ABR in **Right** ear, Normal ABR in **Left** ear

27


Otoacoustic Emissions Present In Both Ears



REMARKS: Earphone: Insert
Transient otoacoustic emissions (TEOAE's) were obtained for frequencies 1.5 through 4.0 K.Hz in the left and right ear. Pt to see Dr. Driscoll.
Copy of audiogram to parents.



Speech Reception Right Awareness 75 dB Left Threshold 10 dB
Phoneme/Word Right / / % @ dB SL Left / 100 % @ 30 dB SL
Recognition: Right / / % @ dB SL Left / / % @ dB SL
Hearing Class: Right D Left A Binaural Hearing Loss, percent: 20.6
Pure tone AC avg: Right 107.5 Left 3.8 Air - Bone Gap: Right NA Left NA

Impittance/Acoustic Reflexes



	500	1K	2K
CONTRA Sound Right			
Reflex (HL)	NR	NR	NR
Decay (sec.)			
Reflex (HL)			
Decay (sec.)			
SOUND Sound Left	500	1K	2K
Reflex (HL)		80	75
Decay (sec.)			
Reflex (HL)			
Decay (sec.)			

Contra-lateral acoustic reflexes present: sound in LEFT ear



Lessons Learned



A profound unilateral hearing loss can be caused by the absence of an auditory nerve and cochlear function could be normal in such cases.

31



Temporal Bone Study of Premature Infants

(Courtesy of Mass. Eye and Ear, published June 2001 Archives Otolaryngology)

Abnormal anatomical findings in autopsies of a group of pre-mature babies

Note missing inner hair cell, normal nerve fiber count inside the habenula perforata, and normal outer hair cells which would lead to normal emissions and NO ABR.





ANSD: C1 - 8 year test results 60 dBA SPL

- CNC words: 60% correct for words & 78% phonemes
- HINT sentences in quiet: 97% correct
- BKB-SIN 4 list average: 10.25 dB SNR for 50% correct
- She performs at the average speech recognition levels as our adults.

35

Summary Points

- Hearing loss is typically a loss of **sensitivity** in hearing levels as demonstrated on an audiogram
- But hearing loss can also be a loss of the ability to make fine **temporal** and **frequency** discriminations
- These factors will reduce the ability of children to “understand” spoken language

36

Case Study X01MK DOB June 06


This child was not screened for hearing at birth and her diagnosis has been delayed.

Audiology- February 2008 – Wearing Spirit3 SP digital hearing aids since November 2007. Not wearing HAs consistently but has begun to wear them since Christmas. She is beginning to say “dada”.

		1kHz	2kHz	4kHz
VRA aided sound field dB HL		35	40	50
VRA inserts	Right	85	80	80
	Left	85	80	85
Unmasked bone conduction		50	70	NR



Beaumont June 08 – ABR LEFT & RIGHT and OAEs under GA ABR tracings for rare vs. condensation clicks only show very large cochlear microphonics of .8 to 1.0 uV in each ear at 100 dB nHL. OAEs are absent on this date. Child has patent grommets in situ R/L today.



MRI February 09

T2 axial imaging shows ventriculomegaly involving the 4th, 3rd and lateral ventricles. Most prominent in the left lateral ventricle. There is no transpendymal

oedema. Significant white matter loss bilaterally, more left than right. **Appearances are suggestive of peri ventricular leukomalacia with almost complete loss of white matter on the left. 7th and 8th nerves and inner ear structures appear to be normal bilaterally.**



Lessons Learned



- **ALL** children being considered for a cochlear implant should have MRI of the brain and auditory nerve cross section to visualize the 4 nerves within the internal auditory canal and verification of brain structures.

40



Lessons Learned



- Hearing loss is typically a loss of sensitivity in hearing levels as demonstrated on an audiogram
- But hearing loss can also be a loss of the ability to make temporal discriminations
- Both factors will reduce the ability of children to “understand” spoken language

41

